



Experimental Review of Inclusive $|V_{ub}|$ Measurements

F Muheim*

School of Physics, University of Edinburgh

We review the status of inclusive $|V_{ub}|$ measurements where many new results have become available recently.

1 Introduction

Precise measurements of the CKM matrix elements $|V_{ub}|$ and $|V_{cb}|$ provide stringent consistency tests of the unitarity triangle. The data on the ratio $|V_{ub}|/|V_{cb}|$ form an annulus in the unitarity plane. This can be tested against the apex of the triangle, determined by the measurements of the CP violating phase $\sin 2\beta$ and the $B^0\bar{B}^0$ mixing parameter Δm_d . The BaBar and Belle experiments have recently measured $\sin 2\beta$ with an error of 0.055 [1] and this precision is expected to improve to 0.02 in a few years time. The precision on $|V_{ub}|$ will soon limit unitarity triangle tests. Hence we aim to measure $|V_{ub}|$ to 10% or better in the near future.

2 Status and Methods

Ten Years ago, CLEO and Argus had established that $|V_{ub}|$ was non-zero [2] by observing an excess of leptons at the endpoint of the momentum spectrum (2.3 - 2.6 GeV/c). However, the accuracy of this result, $|V_{ub}|/|V_{cb}| = 0.08 \pm 0.02$, was theoretically limited by the extrapolation into the full phase which were based on (exclusive) models. Since then huge progress has been made by both, theory and experiment. As reported at this workshop [3], Heavy quark effective theory (HQET) and Operator production expansion (OPE) allow a calculation of the inclusive charmless semileptonic decay rate of B mesons, $B \rightarrow X_u \ell \nu$. The distributions for several kinematic variables, e.g. the lepton momentum, p_ℓ , have also been calculated up to order α_S [4]. The theoretical uncertainties in the determination of $|V_{ub}|$ on the b -quark mass, m_b , and the average kinetic energy of the b -quark within the B meson, λ_1 , are now quantifiable.

The experimental challenge is to suppress the background from $B \rightarrow X_c \ell \nu$ transitions. These have a rate that is higher by two orders of magnitude. Only $\sim 10\%$ of $B \rightarrow X_u \ell \nu$ decays have a lepton momentum larger than 2.3 GeV/c and are above the endpoint of the $B \rightarrow X_c \ell \nu$ background. With the large statistics of B mesons available at the B factories, additional methods with lower efficiency but higher purity are now accessible.

An additional kinematic variable has been already employed by the LEP experiments, namely the mass m_X of the hadronic system X_u recoiling against the lepton pair in the rest frame of the B meson. About 70% of the $B \rightarrow X_u \ell \nu$ decays have $m_X < m_D$ where the mass of the D meson, m_D , is the minimum mass m_X for $B \rightarrow X_c \ell \nu$ decays. In $\sim 25\%$ of $B \rightarrow X_u \ell \nu$ decays the invariant mass of the lepton-neutrino pair, q^2 , exceeds the maximum possible value of background $B \rightarrow X_c \ell \nu$ decays. The energy of the hadronic recoil system could also be used to separate signal from background. The distributions of these kinematic variables have different dependencies on m_b and λ_1 . Thus it is important to measure all kinematic distributions.

Here, we give an overview of six new results on inclusive $|V_{ub}|$ measurements which have been made public by the CLEO, BaBar and Belle collaborations during the last year. For details on these measurements the reader is referred to other presentations at this workshop [5],[6]. Measured inclusive branching ratios $B \rightarrow X_u \ell \nu$ are converted into $|V_{ub}|$ measurements using the relation

$$|V_{ub}| = 0.00445 \sqrt{\frac{\mathcal{B}(BX_u \ell \nu)}{0.002} \frac{1.55 \text{ ps}}{\tau_B}} (1 \pm 0.020 \pm 0.052)$$

or equivalent [7],[8],[9]. The first error arises from the perturbative scale dependence and the second error is due to the b quark mass for which an uncertainty of 90 MeV has been assigned. In what follows, we will refer to these two uncertainties as theoretical error.

3 Lepton Endpoint Spectrum

A new measurement on the excess of leptons in the endpoint region has been published by CLEO [10]. The full data set of 9.13 fb^{-1} at the $\Upsilon(4S)$ resonance and 4.35 fb^{-1} just below has been used. They have carefully designed the suppression of the background from non- B decays (continuum) in order to minimise model dependence on the selection cuts. In Fig. 1 we show the resulting lepton momentum spectrum. A signal interval of 2.2 - 2.6 GeV/c was chosen as this gives the smallest total error. A clear excess of $1901 \pm 122 \pm 256$ events, attributed to $B \rightarrow X_u \ell \nu$, is visible in this region. A partial branching ratio of

$$\Delta \mathcal{B}(B \rightarrow X_u \ell \nu) = (0.230 \pm 0.015 \pm 0.035) \times 10^{-3}$$

*Email: F.Muheim@ed.ac.uk




Figure 1. (a) CLEO lepton spectra for on- (points) and scaled off-resonance data (shaded histogram). The unshaded histogram is the sum of the scaled off-resonance and B -decay background. (b) Background subtracted and efficiency corrected lepton spectrum for $B \rightarrow X_u \ell \nu$ (points) and simulation (histogram).

is measured. The systematic error is dominated by the uncertainty in the $B \rightarrow X_c \ell \nu$ background subtraction. CLEO uses their own measurement of the $B \rightarrow X_s \gamma$ energy spectrum [11] to determine the universal shape function of the heavy-light quark pair system [3],[4]. This in turn allows to calculate the fraction of $B \rightarrow X_u \ell \nu$ signal in the above interval, $f_u = 0.130 \pm 0.024 \pm 0.015$, and to extract a $B \rightarrow X_u \ell \nu$ branching ratio. The branching ratio measurement is then converted into a $|V_{ub}|$ result and CLEO obtains

$$|V_{ub}| = (4.08 \pm 0.34 \pm 0.44 \pm 0.16 \pm 0.24) \times 10^{-3}$$

where the errors are separated into experimental, signal fraction (f_u), theoretical and shape function components, respectively.

Using the lepton endpoint method, The BaBar experiment has presented a preliminary measurement based on 20.6 fb^{-1} of data at the $\Upsilon(4S)$ resonance and 2.6 fb^{-1} below [12]. An excess of $1696 \pm 133 \pm 153$ events is observed in the lepton momentum range of 2.3 - 2.6 GeV/c. BaBar measures a partial branching ratio of

$$\Delta \mathcal{B}(B \rightarrow X_u \ell \nu) = (0.152 \pm 0.014 \pm 0.014) \times 10^{-3}$$

where the experimental systematic error is dominated by the uncertainty in the continuum background subtraction. The CLEO measurement of $f_u = 0.074 \pm 0.014 \pm 0.009$ for this momentum interval is used to extrapolate to the full phase space. Due to the larger extrapolation the uncertainty is comparable to the experimental systematic error. From

the $B \rightarrow X_u \ell \nu$ branching ratio BaBar measures a preliminary $|V_{ub}|$ result of

$$|V_{ub}| = (4.43 \pm 0.39 \pm 0.50 \pm 0.25 \pm 0.35) \times 10^{-3}$$

where the errors are separated into experimental, signal fraction, theoretical and shape function components, respectively.

4 Hadronic Recoil Mass

A measurement of $|V_{ub}|$ by reconstructing the mass m_X of the hadronic system recoiling against the lepton pair in the $B \rightarrow X \ell \nu$ decay has been carried out by BaBar [5],[13]. A data sample of about 82 fb^{-1} is used. The large statistics of B meson samples produced at the B-factories permits the use of new methods which have low efficiency, but allow to exploit the separation power of m_X . The experimental task is to separate the spatially overlapping decay products of the two B mesons produced at the $\Upsilon(4S)$. The principle is to fully reconstruct one B meson in many hadronic modes: $B \rightarrow D^{(*)}\pi$, $D^{(*)}\pi\pi^0$, $D^{(*)}3\pi$, ... The hadronic B meson tags the b -quark flavour, and more importantly removes the background from one of the B mesons in the event. The method yields about 4000 B mesons per fb^{-1} . All the remaining particles that are detected in a tagged event must originate from the other B meson. Events are selected if a lepton with a momentum $p_\ell > 1 \text{ GeV}/c$ in the B rest frame is identified among the particles recoiling against the tagging B meson. The energy substituted mass spectrum, m_{ES} , of the tagging B candidates is shown in Fig. 2. The signal to background ratio is large and a fit is used to remove the continuum background. Additional criteria, e.g. charge correlation and a missing mass consistent with a neutrino are applied to increase the purity of this sample. The hadronic recoil mass, m_X , is reconstructed; a constraint fit imposing zero missing mass is employed to improve the m_X resolution. The $B \rightarrow X_u \ell \nu$ signal region is also enriched by vetoing semileptonic B candidates containing $K^\pm$ or K_S^0 decay products. The depleted region is used as a control sample. In Fig. 3a) we plot the measured m_X spectrum. A binned fit is overlaid. The signal region is $m_X < 1.55 \text{ GeV}$ which gives the smallest total error. It contains one bin to minimise the model dependence. An excess of 167 ± 21 events is observed and attributed to $B \rightarrow X_u \ell \nu$. The signal to background ratio is about 2:1. The background subtracted m_X spectrum with finer bins in the signal region is shown in Fig. 3b). To reduce systematic errors, the ratio $R_u = \mathcal{B}(B \rightarrow X_u \ell \nu)/\mathcal{B}(B \rightarrow X \ell \nu)$ is measured. BaBar obtains a preliminary result of

$$\frac{\mathcal{B}(B \rightarrow X_u \ell \nu)}{\mathcal{B}(B \rightarrow X \ell \nu)} = (1.97 \pm 0.27 \pm 0.23 \pm 0.34) \times 10^{-2}$$

where the errors are statistical, systematic and extrapolation into the region above the m_X cut. Using this branching




Figure 2. BABAR m_{ES} distribution for hadronic tagging B mesons in the lepton sample.




Figure 3. (a) BABAR m_X spectrum (dots) and simulation (histograms). (b) Background subtracted data (dots) and $B \rightarrow X_u \ell \nu$ simulation overlaid.

ratio measurement BABAR presents the following preliminary $|V_{ub}|$ result

$$|V_{ub}| = (4.52 \pm 0.31 \pm 0.27 \pm 0.30 \pm 0.26) \times 10^{-3}$$

where the errors are statistical, systematic, m_X extrapolation and theoretical. The extrapolation error still dominates, but improvements in the knowledge of $\bar{\Lambda}$ and λ_1 are expected from measurements in the next few years. The potential of this method is its small systematic error, thus more statistics will allow to improve the precision on $|V_{ub}|$.

Employing a method similar to BABAR, the Belle experiment has presented a preliminary $|V_{ub}|$ result [6], [17]. Fully reconstructed $B^0 \rightarrow D^{*-} \ell^+ \nu$ decays are used as tagging B mesons. Hence, in selected events, both mesons decay semileptonically. The tag removes the decay products of one of the B mesons and substantially suppresses




Figure 4. Background subtracted m_X spectrum (squares) from Belle. The $B \rightarrow X_u \ell \nu$ simulation (histograms) is overlaid.

the continuum background from non- B decays. Since there are two neutrinos in the event, the B momentum cannot be determined from the tagging B meson. The direction of the two B momenta lay on two cones which can be reconstructed up to an azimuthal angle. After mirroring one of the cones on the origin, the B direction must then lay on the cross line of the two cones and can be reconstructed. As with hadronic tags, the hadronic recoil mass m_X is used to separate $B \rightarrow X_u \ell \nu$ decays from $B \rightarrow X \ell \nu$ decays. In the signal region, defined as $m_X < 1.5$ GeV, Belle observes an excess of 172 ± 28 leptons with momenta $p_\ell > 1$ GeV/c as is shown in Fig. 4. This excess correspond to a branching ratio

$$\mathcal{B}(B \rightarrow X_u \ell \nu) = (2.62 \pm 0.63 \pm 0.23 \pm 0.41) \times 10^{-3}$$

where the errors are statistical, systematic and extrapolation into the region above the m_X cut. Belle obtains a preliminary result for $|V_{ub}|$ of

$$|V_{ub}| = (5.00 \pm 0.60 \pm 0.23 \pm 39 \pm 0.36) \times 10^{-3}$$

where the errors are separated into statistical, experimental, m_X extrapolation and theoretical components, respectively.

5 Neutrino Reconstruction

The neutrino reconstruction technique relies on exploiting the hermeticity of the detector to infer the neutrino momentum from the missing momentum. The missing energy is

also measured and allows to require that the missing mass-squared of the event be consistent with zero. For experiments at the $\Upsilon(4S)$ resonance, this method has been pioneered for exclusive $|V_{ub}|$ and $D_s^+ \rightarrow \mu^+ \nu$ measurements [14],[15].




Figure 5. Fit projection for the q^2 spectrum from CLEO (points). The $B \rightarrow X_u \ell \nu$ signal, $B \rightarrow X_c \ell \nu$ and other background simulations are shown as histograms.

Using neutrino reconstruction, the CLEO experiment has presented a preliminary result on $|V_{ub}|$ using events with inclusive charged leptons [16]. The measurement of the neutrino momentum allows the use of kinematic variables in addition to the charged lepton energy, E_ℓ . The hadronic recoil mass squared, m_X^2 , the invariant mass of the lepton pair, q^2 , and $\cos \theta_{W\ell}$, the helicity angle of the virtual W are used here. Selected events are required to have a lepton with momentum $p_\ell > 1$ GeV/c. A simultaneous fit to the quantities q^2 , m_X^2 , and E_ℓ is performed. The $B \rightarrow X_u \ell \nu$ signal region is restricted to the range $q^2 > 10$ GeV² and $m_X^2 < 2.25$ GeV². In Fig. 5 we show the q^2 projection of the fit with the m_X cut applied for a data sample of 9.4 fb⁻¹. From this fit CLEO extracts a preliminary result of

$$|V_{ub}| = (4.05 \pm 0.18 \pm 0.58 \pm 0.33 \pm 0.56) \times 10^{-3}$$

where the errors are statistical, experimental, m_X and q^2 extrapolation and theoretical.

The Belle experiment has also presented a preliminary $|V_{ub}|$ result using events with charged leptons and neutrino reconstruction [6],[17]. In addition, the particles in each event are separated into a tagging and a signal B meson by employing a combinatorial annealing [18]. The annealing procedure minimizes the likelihood ratio of random over the correct assignments of particles to the two




Figure 6. Background subtracted q^2 spectrum (points) from Belle. The $B \rightarrow X_u \ell \nu$ simulation (histograms) is overlaid.

B mesons in the events. The efficiency of this method is about 0.3%. A control sample of $B^0 \rightarrow D^{*-} \ell^+ \nu$ decays is used to check the annealing and to correct for efficiency differences between data and simulation. After annealing, all kinematic variables of interest are available, for example q^2 and the hadronic recoil mass m_X in the B mesons decaying semileptonically. The signal is extracted in the region of $q^2 > 7$ GeV² and $m_X < 1.5$ GeV. Belle measures an excess of 1148 events over background is measured with a signal to background ratio of 0.27. In Fig. 6 we plot the background subtracted q^2 spectrum with the m_X cut applied. Also shown are $B \rightarrow X_u \ell \nu$ simulations for two values of m_b . A preliminary branching ratio of

$$\mathcal{B}(B \rightarrow X_u \ell \nu) = (1.64 \pm 0.14 \pm 0.36 \pm 0.36) \times 10^{-3}$$

is measured. The errors are statistical, systematic and extrapolation into the region above the m_X and q^2 cut. From this measurement, Belle obtains a preliminary $|V_{ub}|$ result of

$$|V_{ub}| = (3.96 \pm 0.17 \pm 0.44 \pm 0.43 \pm 0.29) \times 10^{-3}$$

where the errors are separated into statistical, experimental, m_X and q^2 extrapolation and theoretical components, respectively.

6 Summary and Conclusions

The B-factories experiments, BaBar and Belle, and CLEO have recently produced many new results for the CKM matrix element $|V_{ub}|$. Different kinematic variables have been used in the results which are based on measurements of




Figure 7. Compilation of inclusive $|V_{ub}|$ results by the Heavy Flavour Averaging Group for the winter conferences 2003.

the inclusive $B \rightarrow X_u \ell \nu$ decay rate. In Fig. 7, we show a compilation of all $|V_{ub}|$ results, carried out by the Heavy Flavour Averaging Group (HFAG) for the winter conferences 2003 [19]. A procedure to average these results is being developed. We have made a preliminary average of all inclusive $|V_{ub}|$ measurements made by the BABAR, Belle and CLEO experiments and obtain

$$|V_{ub}| = (4.32 \pm 0.57) \times 10^{-3}.$$

Statistical and detector errors are treated uncorrelated whereas the extrapolation and theoretical errors are correlated and rescaled. A arbitrary 5% error is assigned to the validity of the shape function being equal for $b \rightarrow s \gamma$ and $B \rightarrow X_u \ell \nu$. Hence we consider this a conservative average.

In this review, we have shown that substantial experimental and theoretical progress has been made over the last decade for measuring $|V_{ub}|$, a crucial parameter in the unitarity triangle. The B-factories allow the use of new experimental methods. The single most precise measurement is BABAR result using the hadronic recoil mass m_X spectrum. The precision of this result is 14% which is better than the average of the LEP results [20]. Excellent neutrino reconstruction is crucial for measuring q^2 which has lower sensitivity to m_b than m_X . The theoretical precision will improve as more precise measurements of the moments of the shape function moments, $\bar{\Lambda}$ and λ_1 , will become available. The aim is to reduce the error on $|V_{ub}|$ to substantially better than 10% over the next five years.

References

1. Gerhard Raven, these proceedings.
2. R. Fulton *et al.* [CLEO Collaboration], Phys. Rev. Lett. **64**, 16 (1990); J. Bartelt *et al.* [CLEO Collaboration], Phys. Rev. Lett. **71**, 4111 (1993); H. Albrecht *et al.* [ARGUS Collaboration], Phys. Lett. B **234**, 409 (1990).
3. M. Luke, these proceedings and references therein.
4. F. De Fazio and M. Neubert, JHEP **9906**, 017 (1999) [arXiv:hep-ph/9905351].
5. A. Sarti, these proceedings.
6. H. Kakuno, these proceedings.
7. N. Uraltsev, Int. J. Mod. Phys. A **14**, 4641 (1999), arXiv:hep-ex/9905520.
8. A.H. Hoang, Z. Ligeti and A.V. Manohar, Phys. Rev. D **59**, 074017 (1999), arXiv:hep-ex/9811239.
9. K. Hagiwara *et al.* [Particle Data Group Collaboration], Phys. Rev. D **66**, 010001 (2002).
10. A. Bornheim *et al.* [CLEO Collaboration], Phys. Rev. Lett. **88**, 231803 (2002).
11. S. Chen *et al.* [CLEO Collaboration], Phys. Rev. Lett. **87**, 251807 (2001).
12. B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0207081.
13. B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0307062.
14. J. P. Alexander *et al.* [CLEO Collaboration], Phys. Rev. Lett. **77**, 5000 (1996).
15. D. Acosta *et al.* [CLEO Collaboration], Phys. Rev. D **49**, 5690 (1994).
16. A. Bornheim *et al.* [CLEO Collaboration], arXiv:hep-ex/0207064.
17. A. Sugiyama, presented at Moriond, March 2003.
18. S. Kirkpatrick *et al.*, Science **220** 4598 (1983).
19. Heavy Flavor Averaging Group, <http://www.slac.stanford.edu/xorg/hfag/>
20. R. Barate *et al.* [ALEPH Collaboration], Eur. Phys. J. C **6**, 555 (1999); M. Acciarri *et al.* [L3 Collaboration], Phys. Lett. B **436**, 174 (1998); P. Abreu *et al.* [DELPHI Collaboration], Phys. Lett. B **478**, 14 (2000); G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C **21**, 399 (2001);